

laser disk decompression

1) introduction

The long term outcome, the complications and suboptimal results which may accompany open disk surgery have led to the early development of other treatment techniques that would avoid a surgical approach through the spinal canal and an extensive disk ablation. Percutaneous removal of nucleus pulposus has been performed with a variety of chemical and mechanical techniques for the past several years. These techniques consist of removing all or part of the nucleus pulposus to induce more rapid healing of the pathological lumbar disk. Percutaneous nucleotomy is now widely used, as it is much less invasive than surgical discectomy. To date, the most promising of these minimally invasive therapies used for treatment of disk herniations is Percutaneous Laser Disc Decompression (PLDD).




Fig 1: Lumbar disk anatomy




Fig 3: Disk herniation




Fig 2: Pathology of disk herniation

2) introduction

Histopathological and bacteriological studies are often needed in musculoskeletal lesions to establish a definitive diagnosis. In such cases Percutaneous Musculoskeletal Biopsy (PMSB) has become a routine procedure.

Advantage of PMSB compared to surgical biopsy :

- No weakening of bone structures through the surgical removal. This is particularly true in weight-bearing bones and in children, thus avoiding immobilization or osteosynthesis.
- No limitation of activity.
- No or minimal soft-tissue injury. No extensive hospitalization, procedure on an out-patient basis.
- No general anesthesia, the PMSB can easily be performed under local anesthesia and neuroleptanalgesia.
- Minimal recovery time No scars Lower cost

3) indications contraindications


Indications

Patient selection is crucial for treatment effectiveness. The indications are : contained disk herniations determined by CT scan or MR imaging with positive and consistent neurologic findings (leg pain of greater intensity than back pain, positive straight-leg-raising test, decreased sensation, normal motor response and tendon reflex) with failure of 6 weeks of conservative therapy


Contraindications

The contraindications are nerve paralysis, hemorrhagic diathesis, spondylolisthesis, spinal stenosis, previous surgery at the same level, significant psychological disorders, significant narrowing of disk space, workplace injuries with monetary gain, local infection of cutaneous or subcutaneous or muscular layers.

4) Technique: sequence of operations

Technique: sequence of operations		
1		Placement
2		Material
3		Dual guidance

4



Local anesthesia

5





5) technique

5 - 1 - placement

The patient is placed in a prone position on the CT table. In order to open up the posterior aspect of the disk space, rolls are positioned under the abdomen to place the lumbar spine in a semiflexed position. This is particularly helpful for the L5-S1 level. The entry point and the pathway are determined by CT, avoiding the nerve root and visceral structures. At the L5-S1 level curved needles are usually necessary.




Fig 9: PLDD CT pathway



Fig 8: PLDD CT disk herniation



Fig 10: PLDD CT pathway

5 - 2 - material

Laser : we use a diode laser (Diomed, Cambridge UK) and a neodymium : yttrium-aluminum-garnet (Nd:YAG) 1064-nanometer wavelength laser (LaserSonics, Milpitas, CA) for PLDD.

- Optical fiber, Y connector
- 22-gauge needle, 18-gauge needle, scalpel
- Iodine, 1% lidocaine Sterile drapes, tampons



Fig 11: Diomed laser (Diomed, Cambridge, UK)



Fig 12 : optical fiber, 22 and 18-g. needle, scalpel, iodine, 1% lidocaine

5 - 3 - guidance

PLDD is performed under dual guidance with a combination of CT and fluoroscopy. Two mobile fluoroscopy monitors are placed in front of the physician along with a CT monitor. At any time the operator can switch from CT to fluoroscopy and vice versa. Once the entry point is determined by CT, a lateral fluoroscopy view is obtained at the desired disk level. In this way, the operator can visualize the pathway and the correct angulation of the needle. In most of the cases the angulation is oblique in the three planes, especially at L5-S1 level.



Fig 13: the patient is placed in prone position. dual guidance CT + fluoroscopy



Fig 15: dual-guidance CT




Fig 14: dual-guidance fluoroscopy

5 - 4 - local anesthesia

The procedure is started respecting strict sterility. The skin subcutaneous layer, lumbar muscles and the articular process are infiltrated by local anesthesia (1% lidocaine) with a 22-gauge needle 9 cm long. The position of the 22-gauge needle is checked by fluoroscopy and CT.




Fig 16: local anesthesia



Fig 18 : CT control of local anesthesia



Fig 17: local anesthesia

5 - 5 - disk puncture

A short scalpel cut is made in the skin. Through the skin incision the 18-gauge needle is inserted under continuous lateral fluoroscopy control parallel to the 22-gauge needle. The tip of the 18-gauge needle must reach the posterior part of the nucleus pulposus. Patient must be monitored for pain during the whole intervention and the needle has to be repositioned if radicular pain occurs. In order to confirm contained disk herniation, or if any doubt persists, a discography can be performed just before PLDD.




Fig 19:18-g.needle placement



Fig 21: scalpel incision of the skin




Fig 20: 18-gauge needle placement under fluoroscopy control



Fig 23: 18-gauge needle placement



Fig 22: 18-gauge needle placement under CT control

5 - 6 - optical fiber placement and check

The optical fiber is checked. Before the optical fiber is placed, it is inserted in an 18-gauge needle mounted by a side-arm fitting to measure the appropriate length of the fiber. The distal part of the optical fiber should extend 5 mm beyond the needle tip. The proper length of the fiber is indicated with sterile strip to avoid excessive advancement of the fiber (placement 1). After removal of the stylet of the 18-gauge needle, the optical fiber is inserted into the disk (placement 2,3,4). The distal part of the optical fiber should extend 5 mm beyond the needle tip .




Fig 24: optical fiber check



Fig 25: optical fiber placement 1



Fig 26: optical fiber placement 2



Fig 27: optical fiber placement 3

5 - 7 - Laser disk ablation

When satisfactory needle position is obtained, the laser procedure begins. The laser is turned on to produce 15-20 watts with 0.5 to 1 second pulses and 4 to 10 second intervals depending on patient comfort. Recommended laser doses for PLDD range from 1200 to 1500 joules for L1-L2, L2-L3, L3-L4 and L5-S1 levels and 1500 to 2000 joules for L4-L5. Using this technique with these short exposure times, there is no heating of the adjacent bone structures. A CT scan is performed every 200 joules at the disk level to visualize the vaporized area. The patient must be able to communicate and respond to pain during the entire procedure. General anesthesia is therefore absolutely contraindicated. If pain occurs, the intervals between pulses are increased and aspiration is applied to reduce pressure within the disk.




Fig 29: laser ablation





Fig 30: CT control of laser ablation

5 - 8 - Follow-up

For 2 weeks after the intervention, some positions that could induce hyperkyphosis as well as athletic activities should be restricted. Resolution of leg pain is usually obtained within one week to two months. The two most critical elements to successful PLDD are proper patient selection and correct needle placement.



Histogram : resolution of leg pain after PLDD MacNab Good. On Yaxis : % of patients with leg pain; on X axis : Days

6) complications

Complications of PLDD under CT and fluoroscopic guidance are rare :

- The major complication of PLDD is septic discitis. To avoid this complication, severe sterility during the intervention is mandatory.
- Thermic aseptic discitis is another complication, leading to severe backache for 3 to 6 months.
- Recurrence of disk herniation or free fragment evacuation with recurrence of leg pain.

7) results

From 1987 to 1997, 169 patients with herniated lumbar disk and radicular pain were treated by PLDD on an out-patient basis. There were 93 male and 76 female patients. The oldest was 71 years and the youngest was 12 years old. Mean age was 42 years. The longest follow-up was 6 years, the average follow-up 19 months . MacNab criteria were used to grade the response to treatment. The overall success rate was 76% according to

MacNab's criteria with 55.6% of GOOD and 20.2% of FAIR. In four cases, the PLDD was performed at two levels. 11 patients with poor results or recurrence were later treated surgically with a satisfying success rate (68%). After 6-12 months, a reduction of disk herniation was observed with CT or MR imaging. These cases were evaluated a second time with a mean follow-up of 53 months, these long-term results were identical. The complications of PLDD under CT and fluoroscopic guidance are rare. The major complication of percutaneous nucleotomy is septic discitis. One patient suffered from a spondylodiscitis. Another suffered for six weeks from severe backache due to an aseptic discitis (three years follow-up). One patient was readmitted 24 hours after PLDD with severe recurrence of leg pain due to free fragment evacuation with upward migration. These data provide encouraging information substantiating the validity of percutaneous laser nucleotomy for contained lumbar disk herniation. The two most critical elements to successful PLDD are : proper patient selection and correct needle placement. Even if further random comparative studies are necessary to confirm these datas, PLDD is a valuable, minimally invasive alternative to conventional surgery for treating disk herniation.

GOOD	Resumed preoperative function. Occasional backache or leg pain. No dependency-inducing medications. Appropriate activity. No objective signs of nerve root damage
FAIR	May be nonproductive if unchanged from preoperative status. Intermittent episodes of mild lumbar radicular pain or low back pain. No dependency-inducing medications. Appropriate activity. No objective signs of nerve root damage
POOR	Subjective : no productivity, continued pain behavior, medication abuse, inactive, compensation litigation focus. Objective : signs of continuing radiculopathy

Fig 1: Mac Nab's criteria.

Location	Cases
L5-S1	87
L4-L5	71
L3-L4	11


Fig 2: location of disk herniation : 169 Cases.

8) cases


Case 1 PLDD for disk herniation with leg pain, satisfactory results of laser vaporization at CT control with air filling the herniation. No complications, good clinical results.




PLDD : CT pathway



PLDD : disk puncture CT control



PLDD : disk puncture fluoroscopy control




PLDD : laser vaporization CT control



PLDD : CT control, laser vaporization gas inside disk herniation (arrow)

Case 2 PLDD with satisfactory results. Note air filling the disk herniation. No complications, good clinical results.




PLDD : disk puncture CT control



PLDD : disk puncture under fluoroscopy control



PLDD : laser vaporization CT control



PLDD : CT control, laser vaporization (red arrow), gas filling annular tear (blue arrow), gas filling disk herniation (yellow arrow)


Case 3 PLDD with good clinical results and no complications.



disk herniation



PLDD : disk puncture CT control



PLDD : disk puncture fluoroscopy control



PLDD : laser vaporization CT control